CHOICES NATIONAL ACTION KIT:



Counseling in WIC Visits to Reduce TV Viewing Strategy Report

CHOICES uses cost-effectiveness analysis to compare the costs and outcomes of different policies and programs promoting improved nutrition or increased physical activity in schools, early care and education and out-of-school settings, communities, and clinics. This strategy report describes the projected national population reach, impact on health and health equity, implementation costs, and cost-effectiveness for an effective strategy to improve child health. This information can help inform decision-making around promoting healthy weight. To explore and compare additional strategies, visit the CHOICES National Action Kit at www.choicesproject.org/actionkit.



TABLE OF CONTENTS

- <u>Page 2</u> **Strategy Profile** | Describes the estimated benefits, activities, resources, and leadership needed to implement the strategy.
- Page 4 National Results | Displays the projected national population reach, impact on health behaviors and prevention of excess weight gain, implementation costs, and cost-effectiveness of the strategy.
- Page 5 Cost Results | Describes the estimated costs by activity and payer needed to implement the strategy nationally.
- Page 7 Health Equity Indicators | Describes the projected impact of implementing the strategy nationally on health equity by race, ethnicity, and income.
- <u>Page 10</u> **Strategy Details & Modeling Methods** | Describes the reach, effect, and cost assumptions used to make national projections for the strategy, and provides links to additional resources related to the strategy.
- <u>Page 12</u> CHOICES National Action Kit: Modeled Outcomes Glossary | Provides definitions for each modeled output displayed in the National Results table.

Page 13 References

SUGGESTED CITATION:

Barrett JL, McCulloch SM, Gortmaker SL, Cradock AL. CHOICES National Action Kit: Counseling in WIC Visits to Reduce TV Viewing Strategy Report. CHOICES Project Team at the Harvard T.H. Chan School of Public Health, Boston, MA; November 2023.

ACKNOWLEDGMENTS:

We thank the following members of the CHOICES Project team for their contributions: Molly Garrone, Banapsha Rahman, Ya Xuan Sun, Amy Bolton, Jenny Reiner, Matt Lee, Zach Ward.

This work is supported by the National Institutes of Health (R01HL146625), The JPB Foundation, the Centers for Disease Control and Prevention (U48DP006376). The findings and conclusions are those of the author(s) and do not necessarily represent the official position of the Centers for Disease Control and Prevention or other funders. The information provided here is intended to be used for educational purposes. Links to other resources and websites are intended to provide additional information aligned with this educational purpose.

Contact the CHOICES Project: choicesproject@hsph.harvard.edu

STRATEGY PROFILE

Describes the estimated benefits, activities, resources, and leadership needed to implement a strategy to improve child health. This information can be useful for planning and prioritization purposes.

Incorporating television time counseling into required Special Supplemental Nutrition program for Women, Infants, and Children (WIC) certification visits among WIC participants with children ages 2-4 through the inclusion of relevant assessment items within the existing screening assessment tools regularly used by WIC clinical staff.

WHAT POPULATION BENEFITS?

Children ages 2-4 who participate in the WIC program.



Relative to not implementing the strategy
Reduce child daily television time which can help promote healthy child weight.



✓ Reduce child daily television time



✓ Prevent cases of obesity



Projected to be cost-saving



 Likely to improve health equity by race, ethnicity, and income

More details available on the CHOICES National Action Kit at choicesproject.org/actionkit

WHAT ACTIVITIES AND RESOURCES ARE NEEDED?

Activities	Resources	Who Leads?
Coordinate planning and delivery of virtual training and educational materials for WIC clinicians and monitor program status	Time for National WIC Coordinator to coordinate and monitor the program	National WIC Coordinator
Develop virtual training and educational materials for WIC clinicians	Time for national WIC agency staff to develop virtual training and educational materials	National WIC Agency Staff
Add relevant assessment items to measure television viewing within tools, tracking, and monitoring systems regularly used by WIC clinicians	Time to update database, tools, and tracking systems	State WIC Agency Information Systems Staff
Train WIC clinicians in using relevant screening tools and motivational interviewing techniques	 Time for state WIC agency staff to prepare for and deliver trainings Time for state WIC clinicians to attend trainings 	State WIC Program Staff

- See our resource library for relevant peer-reviewed publications, research reports, & briefs at choicesproject.org/resource-library
- Learn more about strategy modifications and CHOICES projections of the strategy Counseling in WIC Visits to Reduce TV Viewing for US states:

Arkansas Oklahoma

• Learn more about the evidence for the strategy Counseling in WIC Visits to Reduce TV Viewing in the CHOICES peer-reviewed publication:

Kenney et al. 2021. Child Obes

Adapted from CHOICES Strategy Profile: Counseling in WIC Visits to Reduce TV Viewing. CHOICES Project Team at the Harvard T.H. Chan School of Public Health, Boston, MA; August 2023.

NATIONAL RESULTS

Projected national population reach, impact on health behaviors and prevention of excess weight gain, implementation costs, and cost-effectiveness of the strategy. These national results may help inform your organization's decision-making around promoting healthy weight.

DESCRIPTION	Incorporating television time counseling into WIC clinical visits among WIC participants with children ages 2-4
-------------	---

ОИТСОМЕ	Mean (95% UI)*
BEHAVIOR CHANGE PER PERSON Change in health behavior per person in the first year	1,470 fewer minutes of television viewed (326; 2,690) per year
COST PER PERSON Average annualized cost per person to implement the strategy over the model period	\$0.02 (\$0.02; \$0.02) <u>See Cost Results</u>
POPULATION REACH Reach over the model period	10,300,000 (9,980,000; 10,600,000)
OBESITY PREVENTED Cases of obesity prevented in the final year	7,550 (1,070; 17,500)
CHILD OBESITY PREVENTED Cases of child obesity prevented in the final year	7,550 (1,070; 17,500)
HEALTH EQUITY IMPACT Impact on obesity-related health equity in the final year	Likely to improve health equity by race, ethnicity, & income See Health Equity Indicators
QUALITY-ADJUSTED LIFE YEARS (QALYS) GAINED Quality-adjusted life years (QALYs) gained (totals over the model period)	2,130 (304; 4,690)
OBESITY YEARS PREVENTED Years with obesity prevented (totals over the model period)	47,600 (6,740; 106,000)
HEALTH CARE COSTS SAVED PER \$1 INVESTED Total health care costs saved per total intervention costs over the model period	\$4.47 (\$0.64; \$9.76) <i>Cost-saving</i>
COST PER QALY GAINED Net cost per quality-adjusted life year (QALY) gained (totals over the model period)	Cost-saving 96% likelihood

Projections for the model period 2022-2031 (10 years, inclusive of the start and end years).

Costs are in 2019 dollars and discounted at 3% annually.

*Results displayed are the mean and 95% uncertainty interval (UI). CHOICES calculates 95% uncertainty intervals by running the model 1,000 times and reporting the range (95% of estimates, centered on the mean) of projected outcomes that account for uncertainty from data sources and population projections.

- ✓ Explore our User Guide for more information about the CHOICES National Action Kit at choicesproject.org/action-kit-user-guide
- ✓ Learn more about CHOICES Methods at choicesproject.org/methods
- ✓ Find definitions of each modeled outcome in the Glossary (p.12) at choicesproject.org/action-kit-glossary

COST RESULTS

Describes the estimated costs by activity and payer needed to implement a strategy to improve child health nationally. This information can be useful for planning and prioritization purposes.

This report includes estimates of the implementation costs of Counseling in WIC Visits to Reduce TV Viewing if implemented in each state in the United States. Costs are estimated from a societal perspective, meaning costs needed to implement the strategy are included regardless of who pays or whether the costs are budgetary or opportunity costs.

Average Annual Strategy Implementation Cost by Activity and Payer				
Activity	Resources	Cost per Person†	Payer	Percent of Total Cost
Coordinate planning and delivery of virtual training and educational materials for WIC clinicians and monitor program status	Time for National WIC Coordinator to coordinate and monitor the program	\$0.01	Federal government	65%
Develop virtual training and educational materials for WIC clinicians	Time for national WIC agency staff to develop virtual training and educational materials	\$0.002	Federal government	11%
Add relevant assessment items to measure television viewing within tools, tracking, and monitoring systems regularly used by WIC clinicians	Time to update database, tools, and tracking systems	<\$0.001	State government	2%
Train WIC clinicians in using relevant screening tools and motivational interviewing techniques	Time for state WIC agency staff to prepare for and deliver trainings Time for state WIC clinicians to attend trainings	\$0.004	Local government, federal government	22%
TOTAL		\$0.02		100%

Costs are in 2019 dollars and discounted at 3% per year. Sums may not equal total due to rounding.

[†]Average annualized cost per person to implement the strategy over the model period 2022-2031 (10 years).

Average Annual Strategy Implementation Cost by Payer and Cost Type			
	Cost per Person†		
Payer	All Costs (% of Total)	Budgetary Costs (% of All Costs by Payer)	Opportunity Costs (% of All Costs by Payer)
Federal government	\$0.01 (76%)	-	\$0.01 (100%)
State government	<\$0.001 (21%)	+	<\$0.001 (100%)
Local government	\$0.004 (2%)	+	\$0.004 (100%)
School district		#	-
School		+	-
Family/Individual		#	-
Industry		+	-
Nonprofit	-	-	-
Health care		-	-
TOTAL	\$0.02 (100%)	\$0.00 (0%)	\$0.02 (100%)

Costs are in 2019 dollars and discounted at 3% per year. Sums may not equal total due to rounding. †Average annualized cost per person to implement the strategy over the model period 2022-2031 (10 years).

DEFINITIONS

All costs include budgetary and opportunity costs.

Budgetary costs refer to the actual financial costs incurred.

Opportunity costs refer to the value of what you have to give up in order to choose something else. For example, if an annual professional development training for bullying prevention is replaced with a training for active physical education, there is no budgetary impact, but costs for teachers to attend the training are considered an opportunity cost. The opportunity cost of their time is included in a cost analysis from a societal perspective.

[→] To compare the costs and impacts of strategies, use the <u>CHOICES National Action Kit comparison builder</u>. The strategy implementation cost tables included in this report may provide information useful for planning purposes.

HEALTH EQUITY INDICATORS

Describes the projected impact of implementing a strategy nationally on health equity by race, ethnicity, and income.

Every person deserves access to healthy foods and drinks and opportunities to be physically active, which can help them grow up and live at a healthy weight. However, obesity levels vary by race, ethnicity, and income. Nationally, current and future projected obesity levels are highest among Black or African American and Hispanic or Latino race/ethnicity groups and populations with low household incomes.¹ These disparities are driven by many forces, including commercial determinants leading to increased intake of highly processed and marketed foods and drinks, as well as structural racism and social and economic determinants of health.²-⁴ Effective policy and programmatic strategies promoting improved nutrition and increased physical activity can reduce health disparities and improve health equity.


KEY TAKEAWAYS

If implemented over 10 years (2022-2031), this strategy is projected to:

- ✓ Prevent 7,550 cases of obesity in 2031
- Prevent cases of obesity in all race and ethnicity groups and groups with lower household income
- ✓ Improve health equity by race, ethnicity, and income

Learn more about CHOICES methods for projecting health equity impacts at <u>choicesproject.org/methods/healthequity</u>

Comparative projected impact of the strategy by race and ethnicity




*All Other Races includes people who identify as American Indian/Alaska Native, Asian, Native Hawaiian or Pacific Islander, Multi-racial, or another race or ethnicity not represented in the categories shown. While each of these groups represent distinct populations with differences in health opportunities, risk, and outcomes, they are summarized together due to limited data in national- and state-level surveillance systems.



The Black or African American and Hispanic or Latino populations are projected to experience preventive benefits that are 3.99 and 5.68 times greater compared to the White population. The comparative impact in each population group compared to the population average is provided in a table on page 9.

Continued on the next page

<u>Comparative projected impact of the strategy by household income as a percentage of the federal poverty level (FPL)</u>





Populations with lower household incomes (185% FPL or less) are projected to experience preventive benefits, while the highest income group (household income >350% FPL) is not projected to experience any preventative benefits because they are not the population focus for this strategy. *The comparative impact in each population group compared to the population average is provided in a table on page 9.*

How is this strategy expected to impact health equity?

Every child deserves opportunities to grow up at a healthy weight. Television viewing can lead to increased risk for excess weight because it exposes children to advertising for unhealthy foods and drinks that influences intake and choice preferences. Further, food and beverage companies disproportionately market less healthy foods and beverages to Black and Hispanic/Latino consumers, and Black and Hispanic/Latino people and people from households with lower incomes spend more time viewing television compared to other population groups. Young children spend an average of two and a half hours per day viewing screens, and children from households with the lowest incomes spend more than half an hour per day viewing broadcast television. One strategy to limit the health risks associated with marketing and advertising is to minimize television viewing time. This can be accomplished through efforts to incorporate screening and counseling about reducing television time into the certification visits required for participants with children ages 2-4 in the Special Supplemental Nutrition program for Women, Infants, and Children (WIC), a program that serves people with lower incomes. This strategy is feasible for states to implement, and it may promote health equity related to healthy weight by reducing television viewing among young children from households with lower incomes.

Projected impact of the strategy by race, ethnicity and income, mean (95% UI)^a

	OBESITY PREVENTED ^b	OBESITY PREVENTED PER 100,000 ^b	COMPARAT	IVE IMPACT ^c
	Cases of obesity prevented in the final year	Cases of obesity prevented per 100,000 people in the final year	Ratio of obesity pre	evented per 100,000
Race and Ethnicity			Compared with White, not Hispanic or Latino	Compared with Population Average
Overall	7,550 (1,070; 17,500)	1.53 (0.22; 3.53)		1.00 (Reference) N/A
Black or African American, not Hispanic or Latino	1,560 (199; 3,750)	2.48 (0.32; 5.93)	3.99 (1.90; 8.48) >99% likelihood of greater impact	1.62 (0.92; 2.29) 96% likelihood of greater impact
Hispanic or Latino	3,550 (439; 8,490)	3.53 (0.44; 8.41)	5.68 (3.43; 11.38) >99% likelihood of greater impact	2.31 (1.76; 2.90) >99% likelihood of greater impact
White, not Hispanic or Latino	1,780 (199; 4,110)	0.62 (0.07; 1.43)	1.00 (Reference) N/A	0.98 (0.35; 1.72) 54% likelihood of lesser impact
All Other Races, not Hispanic or Latino ^d	656 (80; 1,670)	1.50 (0.18; 3.82)	2.41 (0.82; 5.91) 96% likelihood of greater impact	0.41 (0.23; 0.55) >99% likelihood of lesser impact
Household Income as a percentage of the federal poverty level (FPL)			Compared with >350% FPL	Compared with Population Average
Overall	7,550 (1,070; 17,500)	1.53 (0.22; 3.53)		1.00 (Reference) N/A
<130% FPL	5,700 (757; 13,100)	4.84 (0.65; 11.10)	Not compared since no impact in reference group	3.17 (2.72; 3.54) >99% likelihood of greater impact
131-185% FPL	1,660 (159; 3,950)	3.23 (0.31; 7.68)		2.11 (1.22; 3.12) 99% likelihood of greater impact
186-350% FPL	199 (0; 598)	0.16 (0; 0.47)		0.10 (0; 0.26) >99% likelihood of lesser impact
>350% FPL	0 (0; 0)	0 (0; 0)		0 (0; 0) >99% likelihood of lesser impact

Projections for the model period 2022-2031 (10 years, inclusive of the start and end years).

^aResults displayed are the mean and 95% uncertainty interval (UI). CHOICES calculates 95% uncertainty intervals by running the model 1,000 times and reporting the range (95% of estimates, centered on the mean) of projected outcomes that account for uncertainty from data sources and population projections.

^bAll cases of obesity prevented are among children, since all people reached by the strategy would still be children in the final model year.

Ratio that compares cases of obesity prevented per 100,000 in each population group with the reference group. When the value is greater than 1.0 for a population group, we project a greater health benefit for that group compared with the reference group. When the value is less than 1.0, we project a lesser health benefit. Note: Ratios are sensitive to extremely high and low rates, so they should be interpreted in the context of the absolute rates, represented by Obesity Prevented per 100,000 here. Results may differ if estimating absolute rates and relative impacts among children only. Likelihood of greater or lesser impact compared with the reference group is estimated based on running the model 1,000 times.

^dAll Other Races includes people who identify as American Indian/Alaska Native, Native Hawaiian or Pacific Islander, Multi-racial, or another race or ethnicity not represented in the categories shown. While each of these groups represent distinct populations with differences in health opportunities, risks, and outcomes, they are summarized together due to limited data in national- and state-level surveillance systems.

STRATEGY DETAILS & MODELING METHODS

Describes the reach, effect, and cost assumptions used to make national projections for the strategy, and provides links to additional resources related to the strategy.

STRATEGY

The CHOICES model estimated nationwide implementation of a strategy to incorporate counseling to reduce television viewing into the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) certification visits. The strategy would include a modification within the existing assessment tool used to personalize nutrition education, referrals, and food package tailoring that would prompt staff to ask caregivers questions during recertification visits and provide relevant educational resources and guidance. The model includes training WIC staff how to incorporate motivational interviewing to decrease home television viewing in WIC clinical visits.^{11,12}

REACH

This strategy would reach children ages 2-4 from households with low income who participate in the WIC program and remain eligible for WIC and do not move for the duration of the program.^{13,14}

This strategy would have a 10-year reach of 10.3 million children if implemented nationwide.

EFFECT

CHOICES estimates that in 2031, 7,550 cases of childhood obesity would be prevented.

The population reached by this intervention would decrease television viewing time by 1,470 minutes per year. We assumed children ages 2-4 years view 0.42 hours per day of television, with children from households with lower incomes viewing more television time compared with children from households with higher incomes. The strategy would decrease daily television viewing time by 13%. To convert change in television hours viewed per day to change in BMI, we used an estimate of the average change in child BMI per one hour reduction per day of television from two randomized trials (-0.33 BMI units/hour). In the convergence of the support of the convergence of the child below the convergence of the convergence

COST

Implementation of this strategy involves the time for the state WIC office staff to develop and implement a videoconference training to all WIC agencies in their state and the time for the VENA support staff to add a screen time field to their online system.

Incorporating counseling to reduce television viewing into the WIC program would incur an average annual cost per child of \$0.02.

CHOICES METHODS

CHOICES uses cost-effectiveness analysis to compare the costs and outcomes of different policies and programs promoting improved nutrition or increased physical activity in schools, early care and education and out-of-school settings, communities, and clinics. Our methods include:

- **Key partner consultation:** Working with key partners & researchers to identify the most promising programs & policies for evaluation
- U.S. population model: Building a computer model of the U.S. population & projecting Body Mass Index (BMI) changes & health outcomes over time
- Systematic reviews & meta-analyses: Synthesizing scientific literature to estimate the likely effects of promising obesity prevention interventions on BMI & physical activity
- **Cost-effectiveness analysis:** Integrating information on the economic costs & health effects of interventions, utilizing a structured & transparent process
- · Health equity lens: Projecting the impact of effective intervention strategies on population health and health equity

Learn more about CHOICES methods at choicesproject.org/methods.

WHY DOES CHOICES USE BMI AS A POPULATION HEALTH INDICATOR?

CHOICES focuses on programs and policies that can help reverse the societal and environmental conditions that drive increases in excess body weight and that emphasize healthy eating, improved physical activity, and reduced screen viewing. Excess body weight is associated with reduced quality of life and increased risk for chronic diseases like diabetes, heart disease, and cancers, ¹⁶ greater healthcare expenditures, ¹⁷ and increased mortality risk. ¹⁸ Obesity is a category of excess weight defined by body mass index (BMI), which is calculated as the ratio of a person's weight (kg) to their height squared (m²). ¹⁹ Obesity is a chronic health condition recognized by the National Institutes of Health, the American Medical Association, Medicare, and Medicaid.

BMI is a useful population health indicator, although it does have limitations. BMI has been shown to be a good measure of individual-level adiposity, correlating highly (r=0.8) with gold standard measures of percent body fat, among adults, children and adolescents and for different gender and racial and ethnic groups.^{20,21} BMI is relatively simple to collect and easy to calculate, and it is used widely in medical and scientific research to measure population health.

However, weight stigma occurs when people are blamed for their weight. Weight stigma can increase a person's risk of engaging in unhealthy eating behaviors and low levels of physical activity and can reduce both the quality of health care a person receives and their utilization of care, all undermining public health.²² CHOICES evaluates the cost-effectiveness of policies and programs aimed at improving nutrition and physical activity environments, promoting related health behaviors, and promoting a healthy weight across all population groups and BMI levels.

For Additional Information

Contact the CHOICES team at choicesproject@hsph.harvard.edu for additional information about model assumptions.

Kenney EL, Mozaffarian RS, Long MW, Barrett JL, Cradock AL, Giles CM, Ward ZJ, Gortmaker SL. Limiting television to reduce childhood obesity: cost-effectiveness of five population strategies. Child Obes. 2021 Oct;17(7):442-448. doi: 10.1089/chi.2021.0016.

For more information about this strategy, see:

Whaley SE, McGregor S, Jiang L, et al. A WIC-based intervention to prevent early childhood overweight. *J Nutr Educ Behav* 2010; 42(3 Suppl):S47–S51. doi:10.1016/j.jneb.2010.02.010

Robinson TN. Reducing children's television viewing to prevent obesity: a randomized controlled trial. *JAMA*. 1999;282(16):1561-1567. doi:10.1001/jama.282.16.1561

CHOICES NATIONAL ACTION KIT: MODELED OUTCOMES GLOSSARY

Provides definitions for each modeled output displayed in the National Results table.

Modeled Output	Definition
BEHAVIOR CHANGE PER PERSON* Change in health behavior per person in the first year	The change in health behavior a person is projected to have after a strategy is put in place. Health behavior changes may include decreases in sugary drink intake, increases in physical activity, decreases in time spent watching TV, or increases in water intake. Behavior change per person is reported when the strategy aims to improve a specific health behavior and data are available to project how much a behavior would improve.
	Averaged across people who actually receive the strategy.
COST PER PERSON Average annualized cost per person to implement the strategy over the model period	The average annualized cost to implement the strategy over the model period (e.g., 10 years) per person reached over the model period. This includes cost by all payers (government, private sector, non-profit, individual/family). See the Cost Results for a breakdown of implementation costs by activity and payer.
	Averaged across people in the intended population of focus where the strategy is adopted (that is, people who are eligible based on age, income, geographic area, and/or participation in the setting or program of focus, and who could potentially receive the strategy based on estimated adoption rates).
POPULATION REACH* Reach over the model period	The number of people reached by the strategy over the model period. Includes all people in the intended population of focus where the strategy is adopted (that is, people who are eligible based on age, income, geographic area, and/or participation in the setting or program of focus, and who could potentially receive the strategy based on estimated adoption rates).
OBESITY PREVENTED* Cases of obesity prevented in the final year	In the final year of the model, the difference in the projected number of people with obesity if the strategy were not put in place and the projected number of people with obesity if the strategy were put in place.
CHILD OBESITY PREVENTED* Cases of child obesity prevented in the final year	In the final year of the model, the difference in the projected number of children with obesity if the strategy were not put in place and the projected number of children with obesity if the strategy were put in place.
HEALTH EQUITY IMPACT* Impact on obesity-related health equity in the final year	The projected impact on differences in obesity levels between population groups defined by race, ethnicity, and by household income. <u>Learn more about our methods for projecting health equity impacts.</u>
QUALITY-ADJUSTED LIFE YEARS (QALYS) GAINED <i>Quality-adjusted life years (QALYs) gained (totals over the model period)</i>	The difference in total number of quality-adjusted life years (QALYs) in the population over the model period if the strategy were not put in place compared with if the strategy were put in place. A QALY is a measure of both the quantity and quality of life. CHOICES estimates the QALYs gained as a measure of how much implementing a strategy to prevent future excess weight gain could improve the quantity and quality of life for a population. See our User Guide for more information about QALYs.
OBESITY YEARS PREVENTED Years with obesity prevented (totals over the model period)	The difference in total number of person-years lived without obesity if the strategy were not put in place compared with if the strategy were put in place. This measure sums up portions of years lived without obesity across all the persons in the model, comparing the result if the strategy were put in place or not.
HEALTH CARE COSTS SAVED PER \$1 INVESTED Total health care costs saved per total intervention costs over the model period	The amount avoided in health care cost related to excess weight for every dollar spent to implement the strategy over the model period. See the Cost Results for a breakdown of implementation costs by activity and payer.
COST PER QALY GAINED Net cost per quality-adjusted life year (QALY) gained (totals over the model period)	The total cost impact to improve population health in terms of quality-adjusted life years gained. Cost per QALY gained is a measure of cost-effectiveness. It includes costs to implement a strategy, cost savings due to efficiencies when implementing a strategy, and health care cost savings related to reductions in excess weight after a strategy is implemented. See our User Guide for more information about QALYs and cost per QALY gained.

All metrics reported for the population over the model period and discounted at 3% per year, unless otherwise noted. Definitions for these modeled outputs are all written assuming that an intervention is implemented.

^{*} Not discounted.

REFERENCES

- Ward ZJ, Bleich SN, Cradock AL, Barrett JL, Giles CM, Flax C, Long MW, Gortmaker SL. Projected U.S. State-Level Prevalence of Adult Obesity and Severe Obesity. N Engl J Med. 2019 Dec 19;381(25):2440-2450.
- Kumanyika SK. A Framework for Increasing Equity Impact in Obesity Prevention. Am J Public Health. 2019 Oct;109(10):1350-1357.
- Bleich SN, Ard JD. COVID-19, Obesity, and Structural Racism: Understanding the Past and Identifying Solutions for the Future. *Cell Metab.* 2021 Feb 2;33(2):234-241.
- Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, Gortmaker SL. The global obesity pandemic: shaped by global drivers and local environments. *Lancet*. 2011 Aug 27;378(9793):804-14.
 Sadeghirad B, Duhaney T, Motaghipisheh S, Campbell NR, Johnston
- Sadeghirad B, Duhaney T, Motaghipisheh S, Campbell NR, Johnston BC. Influence of unhealthy food and beverage marketing on children's dietary intake and preference: a systematic review and metaanalysis of randomized trials. Obes Rev. 2016 Oct;17(10):945-59. doi: 10.1111/obr.12445. Epub 2016 Jul 18. Erratum in: Obes Rev. 2020 Feb;21(2):e12984.
- Russell SJ, Croker H, Viner RM. The effect of screen advertising on children's dietary intake: A systematic review and meta-analysis. Obesity Reviews. 2019;20(4):554-568.
- Harris JL, Frazier W, Kumanyika S, Ramirez AG. Increasing Disparities in Unhealthy Food Advertising Targeted to Black and Hispanic Youth, Rudd Report. January 2019. https://uconnruddcenter.org/wp-content/uploads/sites/2909/2020/09/TargetedMarketingReport2019.pdf
- Rideout V. The Common Sense census: Media use by kids age zero to eight. San Francisco, CA: Common Sense Media; 2017. Accessed December 14, 2020. https://www.commonsensemedia.org/research/the-common-sense-census-media-use-by-kids-age-zero-to-eight-2017
- Rideout V, Robb MB. The Common Sense census: Media use by kids age zero to eight, 2020. San Francisco, CA: Common Sense Media; 2020. Accessed December 8, 2022. https://www.commonsensemedia.org/research/the-common-sense-census-media-use-by-kids-age-zero-to-eight-2020
- CHOICES Stories from the Field: Oklahoma Takes Action to Improve Child Health. Oklahoma State Department of Health & Oklahoma Department of Human Services, Oklahoma City, OK, and the CHOICES Project Team at the Harvard T.H. Chan School of Public Health, Boston, MA; May 2020. Available at: https://choicesproject.org/publications/stft-oklahoma-improve-child-health
- Kenney EL, Mozaffarian RS, Long MW, Barrett JL, Cradock AL, Giles CM, Ward ZJ, Gortmaker SL. Limiting Television to Reduce Childhood Obesity: Cost-Effectiveness of Five Population Strategies. *Child Obes*. 2021. doi: 10.1089/chi.2021.0016.
- Whaley SE, McGregor S, Jiang L, et al. A WIC-based intervention to prevent early childhood overweight. *J Nutr Educ Behav* 2010; 42(3 Suppl):S47–S51. doi: 10.1016/j.jneb.2010.02.010.
 Thorn B, Tadler C, Huret N, Ayo E, Trippe C. WIC Participant and
- Thorn B, Tadler C, Huret N, Ayo E, Trippe C. WIC Participant and Program Characteristics 2014. US Dept of Agriculture, Food and Nutrition Services. 2015:363. Accessed November 30, 2021. Available at: https://fns-prod.azureedge.net/sites/default/files/ops/WICPC2014. pdf
- Epstein LH, Roemmich JN, Robinson JL, et al. A randomized trial of the effects of reducing television viewing and computer use on body mass index in young children. Arch Pediatr Adolesc Med. 2008;162(3):239-245. doi:10.1001/archpediatrics.2007.
- Robinson TN. Reducing children's television viewing to prevent obesity: a randomized controlled trial. JAMA. 1999;282(16):1561-1567. doi:10.1001/jama.282.16.1561.
- Centers for Disease Control and Prevention. Consequences of Obesity. Accessed September 13, 2023 at: https://www.cdc.gov/obesity/basics/consequences.html.
- Ward ZJ, Bleich SN, Long MW, Gortmaker SL. Association of body mass index with health care expenditures in the United States by age and sex. PLoS ONE. 2021 Mar;16(3): e0247307. doi10.1371/journal. pone.0247307
- Ward ZJ, Willett WC, Hu FB, Pacheco LS, Long MW, Gortmaker SL. Excess mortality associated with elevated body weight in the USA by state and demographic subgroup: A modelling study. eClinicalMedicine. 2022 Apr;48. doi:10.1016/j.eclinm.2022.010142
- eClinicalMedicine. 2022 Apr;48. doi:10.1016/j.eclinm.2022.101429

 19. Centers for Disease Control and Prevention. Obesity Basics. Accessed September 13, 2023 at: https://www.cdc.gov/obesity/basics/index.html
- Woolcott OO, Bergman RN. Relative fat mass (RFM) as a new estimator of whole-body fat percentage – A cross-sectional study in American adult individuals. Sci Rep. 2018 Jul 20;8(1):10980.
- Woolcott OO, Bergman RN. Relative Fat Mass as an estimator of wholebody fat percentage among children and adolescents: A cross-sectional study using NHANES. Sci Rep. 2019 Oct 24;9(1):15279.
- Puhl RM, Heuer CA. Obesity stigma: Important considerations for public health. Am J Public Health. 2010;100(6):1019-1028. doi. org/10.2105/AJPH.2009.159491