

Executive Summary

Sugary drink consumption has been linked to excess weight gain, obesity, incidence of type 2 diabetes, heart disease, and cancer. Federal, state, and local governments have considered implementing excise taxes on sugary drinks to reduce consumption, prevent obesity, and provide a new source of government revenue.¹⁻³ In Massachusetts, legislative measures to introduce a tiered sugary drink excise tax have been proposed.^{4,5} The most recent proposed bills specify that tax revenue be dedicated to benefits, services, and programs, including universal free school meals and provision of healthy meals in Head Start and other high need early education settings, for communities most impacted by health inequity and burdened by chronic health outcomes related to sugary drink consumption.^{4,5}

We modeled implementation of a state excise tax on sugary drinks in Massachusetts. Consistent with current policy proposals, we assumed tiered tax rates depending on the sugar content of the beverage: \$0.01/ounce for beverages with more than 7.5 but less than 30 grams of sugar per 12 fluid ounces and \$0.02 for beverages with more than 30 grams of sugar per 12 fluid ounces. CHOICES cost-effectiveness analysis compared the costs and outcomes of implementing a tax with the costs and outcomes expected if the tax were not implemented over 10 years (2023-2032).

The sugary drink excise tax on distributors is projected to be cost-saving. This means that the tax would save more in future health care costs than it costs to implement. This is without consideration of the potential revenue that would be generated, where a tiered \$0.01-\$0.02/ounce statewide excise tax on sugary drinks in Massachusetts could raise as much as \$226 million to \$322 million in annual revenue. Among Massachusetts residents, the tax is projected to decrease sugary drink consumption, prevent over 62,000 of cases of obesity, and save \$937 million in health care costs. People who consume sugary drinks are projected to spend less on these drinks with the excise tax in place. We also project that Black and Hispanic/Latinx Massachusetts residents will experience a greater than average reduction in obesity levels after the tax is implemented, leading to improved health equity. These results are summarized below and in the complete report. Projected results for a \$0.02/ounce state excise tax based on the volume of sugary drinks were similar.

Health impact of a \$0.01-\$0.02/ounce state excise tax on sugary drinks



108 FEWER 12-OZ SUGARY DRINKS SERVINGS PER PERSON

in the first year



62,500 CASES OF OBESITY PREVENTED

in 2032



2,450 CASES OF DIABETES PREVENTED



119,000 FEWER DECAYED, MISSING, OR FILLED TEETH

among Medicaid participants

Cost impact of a \$0.01-\$0.02/ounce state excise tax on sugary drinks







in the first year



DENTAL DECAY TREATMENT COST SAVINGS

for Medicaid

Suggested Citation: McCulloch SM, Barrett JL, Reiner JF, Cradock AL, Gortmaker SL. Massachusetts: Sugary Drink Excise Tax. The CHOICES Learning Collaborative Partnership at the Harvard T.H. Chan School of Public Health, Boston, MA; February 2024. For more information, please visit www.choicesproject.org.

The design for this report and its graphics were developed by Molly Garrone, MA and partners at Burness.

This work is supported by The JPB Foundation and the Centers for Disease Control and Prevention (U48DP006376), and the National Institutes for Health (R01HL146625). The findings and conclusions are those of the author(s) and do not necessarily represent the official position of the Centers for Disease Control and Prevention or other funders.

\$0.01-\$0.02/OUNCE STATE EXCISE TAX ON SUGARY DRINKS

Background

Overconsumption of added sugars is common, with more than half of the United States population ages two years and older exceeding the daily recommended limit for added sugars put forth in the 2015-2020 Dietary Guidelines for Americans. Sugary drinks (defined as all drinks with added caloric sweeteners) are the number one source of added sugars that Americans consume.

In 2018, the beverage industry spent \$1 billion to advertise sugary drinks in television, digital platforms (internet and mobile), radio, magazines, newspapers, coupons, and outdoor venues in order to drive preferences and purchases of sugary beverages. Beverage companies frequently target their sugary drink

In this report, Black or African American, White, and All Other race and ethnicity groups refer to people of non-Hispanic/ Latinx ethnicity.

All Other Races includes people of Asian, Native Hawaiian and Pacific Islander, American Indian and Alaska Native, and two or more races.

advertising towards youth, and are more likely to target Black and Hispanic/Latinx youth. Additionally, Black and Hispanic/Latinx populations are less likely to be the audience for marketing of healthy drinks like water.8 According to recent estimates, approximately 65% of adolescents in Massachusetts report consuming soda regularly, and 11% consume at least one serving of soda per day.9 Higher than average sugary drink consumption levels are common among Hispanic/Latinx and Black Massachusetts youth.9

Strong evidence links increased consumption of sugary drinks to higher risk for obesity and other diseases that are tied to what people eat, such as type 2 diabetes,^{10,11} and the prevalence of these diseases are higher among people with lower income and Black and Hispanic/Latinx Massachusetts residents.^{12,13} An estimated 36% of adults¹³ and 17% of youth¹⁴ in Massachusetts have obesity. If current trends continue, 42% of adults in Massachusetts will have obesity by 2030.¹³

Taxes have emerged as a promising strategy to reduce consumption of sugary drinks. In addition to the potential of a sugary drink tax to reduce obesity, it has cost implications as well. This report models the projected effect of a sugary drink excise tax on health, disease outcomes, and health care cost savings over the next decade.

Projected Impact of a Sugary Drink Excise Tax in Massachusetts

We modeled implementation of a Massachusetts excise tax on sugary drinks, at a tax rate depending on the sugar content of the beverage (i.e., a tiered tax): \$0.01/ounce for beverages with more than 7.5 but less than 30 grams of sugar per 12 fluid ounces and \$0.02 for beverages with more than 30 grams of sugar per 12 fluid ounces. All drinks with added caloric sweeteners were considered to be taxed, while 100% juice, milk products, and beverages with 7.5 or fewer grams of sugar per 12 fluid ounces were considered exempt. As an alternative scenario, we modeled implementation of a \$0.02/ounce state excise tax based on the volume of sugary drinks (i.e., a flat tax).

5

Key Terms

Cost-saving: saves more in future health care costs than it costs to implement (not considering the potential tax revenues)

Excise tax: a consumption tax collected from retailers or distributors; it can be reflected in the posted price (a sales tax in contrast is applied after purchase of the item)

Pass-through rate: how much of the excise tax on retailers or distributors is passed on to consumers as an increase in shelf price; a percent ranging from 0% (none of the tax) to 100% (all the tax), or even greater than 100% (more than the amount of the tax)

Price elasticity of demand: how much consumer purchasing behavior changes following a change in price of an item

Results: What Did We Find?

We project that implementation of a state excise tax on sugary drinks, at a tax rate of \$0.01/ounce or \$0.02/ounce depending on sugar content, has greater than a 99% likelihood of being cost-saving by saving more in future health care costs than it would cost to implement. Among the Massachusetts population, it would prevent 62,500 of cases of childhood and adult obesity, prevent new cases of diabetes, increase healthy life years, prevent tooth decay, improve health equity, and save more in future health care costs than it will cost to implement. Projected results for a \$0.02/ounce state excise tax based on the volume of sugary drinks were similar. For complete results, see the Appendix beginning on page 16.

Model results are presented as the most likely estimate as well as a 95% uncertainty interval (95% UI). The 95% UI can be thought of as a likely range that is estimated by considering uncertainty from data sources and population projections and calculating a central range in which 95 percent of the model results fell.



How many people would be affected by the tax?

This can be thought of as reach or the number of people affected by the strategy. Based on our modeling, the table below presents the estimated number of people affected by the tax in the first year and the number of people affected by the tax over 10 years.

Outcome	Number of people affected by the tax	95% UI
First Year Population Reach*	6.57 million	(6.55 million; 6.60 million)
10-Year Population Reach*	7.18 million	(7.14 million; 7.23 million)

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.

*Not discounted.

What effect would the tax have on sugary drink consumption and spending?

Economic studies indicate that with a sugary drink tax on distributors, consumers will buy less of these products. Assuming fewer purchases result in lower consumption, a \$0.01-\$0.02/ounce tax will decrease the intake of sugary drinks. Compared to projections of sugary drink consumption without a tax, the tax is projected to result in lower levels of sugary drink consumption. In Massachusetts, spending on sugary drinks without a tax is estimated to be \$1.56 billion per year. We project a 96% likelihood that individuals and households who purchase sugary drinks will spend less on sugary drinks after the tax is implemented.


Outcome	Impact of the tax on sugary drink consumption & spending	95% UI
Decrease in 12-oz Servings of Sugary Drinks per Person in the First Year*	108	(66; 209)
Decrease in Spending on Sugary Drinks in the First Year per Person Consuming Sugary Drinks*	^{\$} 65	(-\$2; \$231) 96% likelihood of decrease in spending
Decrease in Spending on Sugary Drinks in the First Year per Household*	\$165	(-\$6; \$582) 96% likelihood of decrease in spending
Decrease in Spending on Sugary Drinks in the First Year Overall in Massachusetts*	\$241 million	(-\$8.66 million; \$849 million) 96% likelihood of decrease in spending

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.


*Not discounted.

Average Annual Pre-Tax Sugary Drink Consumption per Person in Massachusetts by Race and Ethnicity



[†]Each serving is 12 ounces. DATA SOURCES: NHANES 2011-2016; Analysis by CHOICES Project, 2020.

Post-Tax Decrease in Sugary Drink Consumption per Person in Massachusetts by Race and Ethnicity*



*In the first year following a sugary drink excise tax of \$0.01-\$0.02/ounce based on sugar content.

†Each serving is 12 ounces.



What effect would the tax have on obesity and related health outcomes, overall and by race and ethnicity?

Compared to projections of obesity and related health outcomes without a tax, the tax is projected to result in fewer cases of obesity and fewer deaths over the 10-year period under consideration. Under the proposed tax, Black Massachusetts residents will experience a preventive health benefit that is 9% greater than the population average, and Hispanic/Latinx Massachusetts residents will experience a preventive health benefit that is 19% greater than the population average.

Outcome	Impact of the tax on obesity and related health outcomes	95% UI
Quality Adjusted Life Years (QALYs) Gained Over 10 Years	29,400	(13,700; 66,500)
Years of Life Gained Over 10 Years	6,350	(2,620; 15,000)
Deaths Prevented Over 10 Years*	1,820	(746; 4,290)
Years with Obesity Prevented Over 10 Years	479,000	(230,000; 1.06 million)
Cases of Obesity Prevented in 2032*	62,500	(29,900; 136,000)
Cases of Childhood Obesity Prevented in 2032*	8,320	(3,390; 17,300)
Cases of Obesity Prevented Per 100,000 in 2032*	978	(470; 2,140)

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.


*Not discounted.

\$0.01-\$0.02/ounce state excise tax on sugary drinks.




in 2032

Pre-Tax Obesity Prevalence in Massachusetts by Race and Ethnicity



 ${\tt DATA\ SOURCES:\ NHANES\ 2011-2016,\ NSCH\ 2003-2018;\ Analysis\ by\ the\ CHOICES\ Project,\ 2023.}$

Comparative Projected Impact of a Sugary Drink Excise Tax in Massachusetts by Race and Ethnicity*



Percentage of the Massachusetts Population by Race and Ethnicity

- 7% Black or African American, not Hispanic/Latinx
- 13% Hispanic/Latinx
- 71% White, not Hispanic/Latinx
- 9% All Other Races, not Hispanic/Latinx

 * With a sugary drink excise tax of \$0.01-\$0.02/ounce based on sugar content.

	Massachusetts: Impact of the tax on behavior and health, by race and ethnicity**			
Outcome	Black or African American, not Hispanic/Latinx	Hispanic/Latinx	White, not Hispanic/Latinx	All Other Races, not Hispanic/ Latinx†
	Mean	Mean	Mean	Mean
	95% UI	95% UI	95% UI	95% UI
Decrease in 12-oz Servings of Sugary Drinks per Person in the First Year*	121 (74; 227)	115 (70; 211)	106 (65; 212)	101 (62; 190)
Quality Adjusted Life	2,060	3,840	21,300	2,200
Years (QALYs) Gained Over 10 Years	(970; 4,460)	(1,850; 8,130)	(9,750; 49,900)	(1,040; 4,720)
Years of Life Gained Over	516	508	4,940	381
10 Years	(155; 1,260)	(153; 1,190)	(2,020; 11,700)	(102; 979)
Years with Obesity Prevented Over 10 Years	35,100 (17,500; 72,600)	77,800 (38,100; 163,000)	330,000 (155,000; 746,000)	36,200 (18,000; 73,600)
Cases of Obesity Prevented in 2032*	4,580 (2,250; 9,470)	10,300 (5,060; 21,000)	42,700 (20,100; 95,200)	4,890 (2,340; 9,830)
Cases of Childhood Obesity Prevented in 2032*	911 (337; 1,940)	2,730 (1,070; 5,700)	3,890 (1,570; 8,000)	792 (314; 1,700)
Cases of Obesity Prevented Per 100,000 in 2032*	1,070 (522; 2,240)	1,170 (573; 2,420)	960 (452; 2,140)	783 (376; 1,590)
Comparison of Cases	1.11	1.21	1.00 (Reference)	0.82
of Obesity Prevented per 100,000 in 2032 vs. White*‡	(0.91; 1.45) 84% likelihood of greater impact	(0.97; 1.68) 94% likelihood of greater impact	N/A	(0.67; 1.06) 94% likelihood of lesser impact
Comparison of Cases	1.09	1.19	0.98	0.80
of Obesity Prevented	(0.93; 1.30)	(1.00; 1.50)	(0.89; 1.04)	(0.69; 0.96)
per 100,000 in 2032 vs. Average*‡	84% likelihood of greater impact	98% likelihood of greater impact	69% likelihood of lesser impact	>99% likelihood of lesser impact

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.

^{*}Not discounted.
**During the modeling period, 2023-2032.

[†]All Other Races includes people of Asian, Native Hawaiian and Pacific Islander, American Indian and Alaska Native, and two or more races.

[‡]The ratio of the projected reduction in obesity prevalence in a particular group compared to the projected reduction in the reference group. A value greater than 1 indicates that the group is projected to see a greater reduction in obesity levels compared to the reference group.



What effect would the tax have on diabetes?

Decreasing the intake of sugary drinks has other health implications. We estimated the impact of the tax on diabetes incidence using the projected estimates of decline in sugary drink consumption. In Massachusetts, the proposed sugary drink excise tax on distributors could lead to a 10% reduction in diabetes incidence after the first two years of implementation.

\$0.01-\$0.02/ounce state excise tax on sugary drinks.



10% REDUCTION IN DIABETES INCIDENCE



2,450 CASES OF DIABETES PREVENTED



What effect would the tax have on tooth decay?

Higher intake of sugar, including in sugary drinks, is associated with increased decayed, missing, and filled teeth (i.e., tooth decay). ^{16,17} We estimated the impact of the tax on tooth decay using the projected declines in sugary drink consumption estimated. In Massachusetts, we estimated that a \$0.01-0.02/ounce tiered tax would lead to 119,000 fewer decayed, missing, and filled teeth among Medicaid recipients and \$9.16 million in savings to Medicaid over 10 years due to a reduction in treatment of tooth decay. For the entire Massachusetts population, we estimated the tax would lead to a reduction of 757,000 decayed, missing, and filled teeth and \$58.2 million in savings for all payers due to a reduction in treatment of tooth decay. A recent study of dental patients indicated that the sugary beverage tax in Philadelphia led to reductions in decayed, missing and filled teeth among both children and adults on Medicaid. These results validate our projections, and also indicate likely improvements in health equity. ¹⁸

\$0.01-\$0.02/ounce state excise tax on sugary drinks...

\$9.16 mill

DENTAL DECAY TREATMENT COST SAVINGS

over 10 years (Medicaid)

\$58.2 mill

DENTAL DECAY TREATMENT TOTAL COST SAVINGS

over 10 years (Societal)



How much would the tax cost to implement?

To implement the proposed policy in Massachusetts, the Massachusetts Department of Revenue would need to process tax statements and conduct audits. Businesses would also need to prepare tax statements and participate in audits, which would require labor from private tax accountants. The Massachusetts Department of Public Health would also lead a communications campaign to promote healthy beverage consumption and health equity. Cost information is based on data from localities with planned or implemented excise taxes on sugary drinks¹⁹ and from programs previously implemented in Massachusetts. The cost and benefit estimates do not include expected tax revenue from distributors (discussed below). Below we include annual and 10-year implementation costs.

Outcome	Costs	95% UI
Annual Implementation Cost	\$380,000	(\$327,000; \$429,000)
Annual Implementation Cost per Person	^{\$} 0.06	(\$0.05; \$0.07)
Total Intervention Implementation Cost Over 10 Years	\$3.80 million	(\$3.27 million; \$4.29 million)

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.



How much would the tax save in health care costs compared to what it costs to implement?

Since we project that the tax saves more in future health care costs than it costs to implement, the tax would be cost-saving. We project a greater than 99% likelihood of cost savings with the tax. The estimated reduction in obesity attributable to the tax leads to lower projected health care costs, offsetting tax implementation costs and resulting in a net cost savings. The difference between total health care costs with no tax implementation and lower health care costs associated with the implementation of a tiered sugary drink tax on distributors represents the health care costs saved; these savings can be compared to the cost of implementing the tax to arrive at the metric of health care costs saved per \$1 invested.

Outcome	Costs	95% UI
Health Care Costs Saved Over 10 Years	\$937 million	(\$397 million; \$2.19 billion)
Net Costs Difference Over 10 Years	-\$933 million	(-\$2.19 billion; -\$393 million)
Health Care Costs Saved per \$1 Invested Over 10 Years	^{\$} 246.39	(\$102.30; \$592.99)

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.



What would the impact of the tax be on Medicaid spending?

Based on the projected reduction in adult obesity prevalence due to the tax (compared to what prevalence would be without the tax), adult obesity-related MassHealth expenditures in Massachusetts are estimated to decrease. This does not include potential reductions in child obesity-related expenditures, and may be an underestimate if adults utilizing MassHealth have higher than average health care costs of obesity.²⁰ A previous analysis found that 10% of adult Medicaid expenditures were due to obesity.²¹ We estimate that, in Massachusetts, obesity accounted for \$1.47 billion of \$18.1 billion total adult MassHealth expenditures in 2021.²² This assumes that 81% of all MassHealth payments are for adults.²² The state paid 22% of total MassHealth expenditures in 2021,²² so we estimate that the state paid \$323 million in adult obesity-related MassHealth expenditures.

†The Medicaid program in Massachusetts is MassHealth.

Outcome	Medicaid spending	95% UI
Reduction in Annual Adult Obesity-Related Medicaid Expenditures (Paid by State)*	\$3.39 million	(\$1.53 million; \$7.89 million)
Reduction in Total Annual Adult Obesity- Related Medicaid Expenditures (Paid by State and Federal)*	\$15.4 million	(\$6.96 million; \$35.8 million)

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.

*Not discounted.



Would the tax be a good value?

We project that the tax would improve health and save more in future health care costs than it costs to implement, making it cost-saving.

The tax would be cost-saving on the basis of the key cost-effectiveness metrics:

- Cost per year with obesity prevented over 10 years
- Cost per quality adjusted life year (QALY) gained over 10 years
- Cost per year of life gained over 10 years
- Cost per death averted over 10 years

Key Considerations for Health Equity

Community members have raised concern that the tax would have a disproportionate economic impact on households with lower incomes. Recent studies of existing sugary drink taxes^{18,23} and conclusions from health economists²⁴ suggest that populations with low income benefit substantially from sugary drink taxes. A recent study from England indicates the tiered tax there resulted in reduced obesity in young girls, with the greatest impacts upon those living in the most deprived areas.²³ Another recent study of dental patients indicated that the sugary beverage tax in Philadelphia led to reductions in decayed, missing and filled teeth among both children and adults on Medicaid.¹⁸



Economic studies indicate that with a sugary drink tax, consumers will buy less of these products.¹⁵ This change in purchasing is substantial, so that consumers can be expected to spend less on sugary drinks after a tax is

implemented. Using sales data from the Rudd Center Revenue Calculator for Sugary Drink Taxes,⁶ we estimate that \$1.56 billion is spent annually on sugary drinks in Massachusetts. With a tiered \$0.01-\$0.02/ounce tax in place, we project that spending on sugary drinks in Massachusetts will decrease by \$241 million in the first year. A typical person in Massachusetts who consumes sugary drinks would spend \$65 less per year on sugary drinks after a tiered \$0.01-\$0.02/ounce tax, and a typical household would spend \$165 less per year. This would free up disposable income for other household purchases. A typical person in Massachusetts who continues to drink sugary drinks after the tax is in place would be expected to pay a tax of about \$1.27 per week, or \$66 per year.

In addition to these changes in spending, reductions in obesity rates due to the tax are projected to be greatest among individuals from households with low income. We also project that there would be greater than average reductions in obesity rates among Black and Hispanic/Latinx Massachusetts residents. Using data on sugary drink consumption in the CHOICES model, the average daily consumption of sugary drinks among people in Massachusetts varies by race and ethnicity group (see pre-tax graphs on pages 4 and 6). Under the proposed tax, Hispanic/Latinx Massachusetts residents would see a 19% greater than average reduction in obesity rates and Black residents would see a 9% greater than average reduction in obesity rates. On that basis, the modeled tax could decrease disparities in obesity outcomes and improve health equity by race and ethnicity.

Implementation Considerations

A tiered \$0.01-\$0.02/ounce statewide excise tax on sugary drinks in Massachusetts could raise as much as \$226 million to \$322 million in annual revenue.⁶ Revenue raised from a sugary drink tax could be invested in communities with lower incomes if the legislature earmarks the tax for this type of use. Cities with current sugary drink taxes in the U.S. have allocated these revenues in a variety of ways, including "increasing access to healthy food and water, educating about nutrition and healthy beverage choices, providing health services, and expanding opportunities for physical activity. Others address social determinants of health such as early childhood education or maintenance of libraries, parks, and recreation centers." Public support for such taxes generally increases with earmarking for preventive health activities.²⁶

There is opposition from the food and beverage industry, which spends billions of dollars promoting their products.²⁷ One concern is the impact on employment. In U.S. cities with sugary drink taxes in place, there is no evidence that the tax has negatively impacted employment.^{28,29} Relatively small beverage excise taxes are currently applied across many states. The proposed tax is likely to be sustainable if implemented based on the successful history of tobacco excise taxes. There is potential for a shift in social norms of sugary drink consumption based on evidence from tobacco control tax and regulatory efforts.³⁰

Modeling Assumptions & Summary of the CHOICES Microsimulation Model



How does an excise tax on distributors affect the price paid by consumers?

An excise tax is incorporated directly into a beverage's shelf price. We assume 100% pass-through of the tax over 10 years and assume the tax rate would be adjusted annually for inflation. Our pass-through rate estimate is supported by empirical studies of excise taxes in Mexico and France that demonstrate near-complete pass-through rates to consumers³¹ and consistent evidence in the US indicating that beverage taxes increase prices, although there is some variation by store type.³²⁻³⁵

The expected change in sugary drink price was estimated based on national sugary drink prices³⁶ and regional sales data for several categories of sugary drinks (i.e., soda, sports drinks, fruit drinks, energy drinks, sweetened teas, sweetened coffees, and enhanced water).⁶ In Massachusetts, we assume the average price of sugary drinks is \$0.068/ounce, and 72% of the taxable sugary drinks that people consume would be taxed at \$0.02/ounce, with 28% taxed at \$0.01/ounce, so a tiered tax would raise prices by about 25%. This means that, for example, the price of a 12-ounce can of soda would increase from \$0.82 to \$1.02/can post-tax.



How does increasing the price of sugary drinks change individual sugary drink consumption?

How much consumers will change their purchases in response to price changes is called price elasticity of demand. We assume for every 10% increase in the price of sugary drinks, there will be a 12% reduction in purchases (a mean own-price elasticity of demand of -1.21). Research on existing sugary drink taxes find price elasticities consistent with this estimate. He Massachusetts, we assume a tiered \$0.01-0.02/ounce tax that raises prices by 25% would reduce purchases by 30%. We assume this 30% reduction in purchases results in a 30% reduction in consumption.

To estimate current sugary drink consumption levels in Massachusetts, we used national estimates of sugary drink consumption by sugar content from NHANES 2011-2016 adjusted to the demographic makeup of the Massachusetts population by age, sex, race, and ethnicity.



What are the health effects of decreasing sugary drink consumption?

Research has shown that decreasing sugary drink consumption can have positive effects on health in youth and adults.



ASSUMPTIONS ABOUT SUGARY DRINKS AND OBESITY RISK

We estimated the impact of a change in sugary drink intake on body mass index (BMI), accounting for dietary compensation, based on rigorous studies identified in evidence reviews.^{19,42} We assumed that people with higher BMI experience greater reductions in excess weight after reducing sugary drink intake, compared with people with lower BMI.⁴³⁻⁴⁵ The relationship among adults was modeled based on the range of estimated effects from four large, multi-year longitudinal studies, which indicated that a one-serving reduction in sugary drinks was associated with an average BMI decrease of 0.21 kg/m² to 0.57 kg/m² in adults over a 3-year period.^{11,46-48} Among youth, we used evidence from a double-blind randomized controlled trial conducted over 18 months, which found that an additional 8-ounce serving of sugary drinks led to a 2.2 pound greater weight gain on average.⁴⁹



ASSUMPTIONS ABOUT SUGARY DRINKS AND DIABETES RISK

We estimated the impact of the tax-induced reduction in sugary drink intake on diabetes incidence for adults ages 18-79 years using a published meta-analysis of the relative risk of developing diabetes due to a one-serving change in sugary drink consumption⁵⁰ as well as state-level estimates of diabetes incidence from the CDC Atlas 2022⁵¹ and prevalence from the Massachusetts Behavioral Risk Factor Surveillance System 2021.¹² On average, each 8.5-ounce serving of sugary drinks per day increases the risk of diabetes by 18%.⁵⁰




ASSUMPTIONS ABOUT SUGARY DRINKS AND TOOTH DECAY

We estimated the impact of a sugary drink excise tax on tooth decay and tooth decay treatment costs using a longitudinal analysis of the relationship between intake of sugars and tooth decay in adults. On average, for every 10 grams higher intake of sugar per day, there is an increase in decayed, missing, and filled teeth (i.e., tooth decay) of approximately 0.10 over 10 years.¹⁶ There are many studies showing a similar relationship between higher intake of sugars and tooth decay in children and youth¹⁷ and thus we assume the same relationship as found in adults. We used the allowable fees for dental services provided to individuals on public insurance set forth in the Code of Massachusetts Regulations (101 CMR 314.00) in 2018⁵² to estimate a Medicaid cost of treating tooth decay as: \$637.83 for a permanent crown and \$119.77 for a filling for children and \$504.83 for a permanent crown and \$86.08 for a filling. These codes reflect treatment for one to four surfaces but do not reflect the actual frequency of multi-surface treatment among Medicaid recipients or for higher reimbursement rates due to temporary crowns or potential flat tax schedules. Using the allowable fees may underestimate the total cost savings of tooth decay treatment projected here as dental providers may charge higher amounts to patients. Based on analysis of data on tooth decay, fillings, and crowns for the U.S. population from NHANES 1988-1994 (the last year crowns and fillings were separately reported),53 we estimate that 78.9% of tooth decay in children and 43.5% of tooth decay in adults is treated. Using this same data set, we estimate that 97% of treatment for children is fillings and 82.5% of treatment for adults is fillings. To estimate Medicaid-specific savings in costs of dental treatment, we used estimates of the number of people enrolled in Medicaid and the proportion of people receiving Medicaid dental services in 2018.54,55

CHOICES Microsimulation Model

The CHOICES microsimulation model was used to calculate the costs and effectiveness of a tax in Massachusetts over 10 years (2023-2032). We forecasted what would happen to a virtual population of residents in Massachusetts with and without a sugary drink tax to model changes in disease and mortality rates and health care costs due to the tax. Cases of obesity prevented were calculated at the end of the model period in 2032. The model was based on peer-reviewed CHOICES methodology, 19,56-58 with updated assumptions and data sources based on new data available and methodological refinements made over time. 59 We created a virtual population of residents in Massachusetts using data from: the U.S. Census, American Community Survey, Behavioral Risk Factor Surveillance System, NHANES, National Survey of Children's Health, 58 the Medical Expenditure Panel Survey, multiple national longitudinal studies, and obesity prevalence data provided by Massachusetts Department of Public Health. Impacts on diabetes, tooth decay, Medicaid spending, and household spending on sugary drinks were calculated based on summary results from the model, not directly via microsimulation.

Of note, the CHOICES microsimulation model does not include annual revenue generation from a state excise tax on sugary drinks in any of the cost-effectiveness calculations. The Rudd Center Revenue Calculator for Sugary Drink Taxes estimates potential annual revenues from excise taxes on sugary drinks and is "intended to provide a rough estimate" for municipalities to consider. The Rudd Center calculator provides revenue estimates for taxes based on beverage volume only, not those based on sugar content like the tax modeled for Massachusetts. Since we assume 72% of taxed sugary drinks consumed fall into the highest sugar content tier, we assume revenue for a \$0.0175/ounce volume tax approximates estimated revenue for a \$0.01-\$0.02/ounce tiered tax. According to the Rudd Center, \$0.0175/ounce excise tax in Massachusetts could raise as much as \$322 million in 2023. Accounting for 10-30% non-compliance as the Rudd Center advises, annual revenue estimates may be \$226 to \$322 million.



References

- 1. American Public Health Association Taxes on Sugar-Sweetened Beverages. 2012.
- Falbe J, Rojas N, Grummon AH, Madsen KA. Higher Retail Prices of Sugar-Sweetened Beverages 3 Months After Implementation of an Excise Tax in Berkeley, California. American Journal of Public Health. 2015;105(11):2194-2201.
- World Health Organization. Global action plan for the prevention and control of noncommunicable diseases 2013–2020 [Internet]. Geneva: WHO. Updated appendix 3, "Best buys" and other recommended interventions for the prevention and control of noncommunicable diseases; [updated 2017; cited 2019 Sep 17]. Available from: https://iris.who.int/bitstream/handle/10665/94384/9789241506236_eng.pdf?sequence=1
- Massachusetts Senate Docket No. 959. An Act to promote healthy alternatives to sugary drinks. https://malegislature.gov/Bills/193/SD959. Filed January 18, 2023. Accessed March 9, 2023.
- Massachusetts House Docket No. 1813. An Act to promote healthy alternatives to sugary drinks. https://malegislature.gov/Bills/193/HD1813. Filed January 18, 2023. Accessed March 9, 2023.
- UCONN Rudd Center. Revenue Calculator for Sugary Drink Taxes. Release: April 13, 2021. http://www.uconnruddcenter.org/revenue-calculator-for-sugary-drink-taxes. Accessed December, 2023.
- U.S. Dept of Health and Human Services & U.S. Dept of Agriculture, 2015-2020 Dietary Guidelines for Americans, 8th Edition, 2015.
- Harris JL, Fleming-Milici F, Kibwana-Jaff A, Phaneuf L. Sugary Drink FACTS 2020: Sugary drink advertising to youth: Continued barrier to public health progress. UConn Rudd Center for Food Policy & Obesity, June 2020. Accessed January 7, 2021 at http://sugarydrinkfacts.org/resources/Sugary%20Drink%20FACTS%202020/Sugary_Drink FACTS Full%20Report final.pdf
- Centers for Disease Control and Prevention. Youth Online. Massachusetts, High School Youth Risk Behavior Survey, 2019. https://nccd.cdc.gov/Youthonline/App/Results.aspx?LID=MA. Accessed November 5, 2020.
- Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. American Journal of Clinical Nutrition. 2013;98(4):1084-1102. 16.
- Nutrition. 2013;98(4):1084-1102. 16.

 11. Chen L, Caballero B, Mitchell DC, et al. Reducing Consumption of Sugar-Sweetened Beverages Is Associated with Reduced Blood Pressure: A Prospective Study among U.S. Adults. Circulation. 2010;121(22):2398-2406.
- Massachusetts Department of Public Health, Office of Data Management and Outcomes Assessment. A Profile of Health Among Massachusetts Adults, 2021. Published January 2023. https://www.mass.gov/doc/a-profile-of-health-among-massachusetts-adults-2021-0/download
- Ward ZJ, Bleich SN, Cradock AL, Barrett JL, Giles CM, Flax CN, Long MW, Gortmaker SL. Projected U.S. State-Level Prevalence of Adult Obesity and Severe Obesity. N Engl J Med. 2019;381:2440-50. doi: 10.1056/NEJMsa1909301
- Massachusetts Department of Public Health. Results from the Body Mass Index Screening in Massachusetts Public School Districts, 2017. https://www.mass.gov/doc/the-statusof-childhood-weight-in-massachusetts-2017/download. Published July 2020. Accessed November 5, 2020.
- Powell LM, Chriqui JF, Khan T, Wada R, Chaloupka FJ. Assessing the Potential Effectiveness of Food and Beverage Taxes and Subsidies for Improving Public Health: A Systematic Review of Prices, Demand and Body Weight Outcomes. Obesity reviews: an official journal of the International Association for the Study of Obesity. 2013;14(2):110-128
- Bernabe E, Vehkalahti MM, Sheiham A, Lundqvist A, Suominen AL. The Shape of the Dose-Response Relationship between Sugars and Caries in Adults. Journal of Dental Research. 2016;95(2):167-172.
- Sheiham A, James WPT. A new understanding of the relationship between sugars, dental caries and fluoride use: implications for limits on sugars consumption. Public Health Nutrition. 2014;17(0):2176-2184.
- Petimar J, Gibson LA, Wolff MS, Mitra N, Corby P, Hettinger G, Gregory EF, Edmondson E, Block JP, Roberto CA. Changes in Dental Outcomes After Implementation of the Philadelphia Beverage Tax. Am J Prev Med. 2023 Feb 28
- Gortmaker SL, Wang YC, Long MW, et al. Three Interventions That Reduce Childhood Obesity Are Projected To Save More Than They Cost To Implement. Health Affairs. 2015;34(11):1932-1939.
- Wang YC, Pamplin J, Long MW, Ward ZJ, Gortmaker SL, Andreyeva T. Severe Obesity In Adults Cost State Medicaid Programs Nearly \$8 Billion In 2013. Health Aff (Millwood). 2015 Nov;34(11):1923-31.
- Trogdon JG, Finkelstein EA, Feagan CW, Cohen JW. State- and payer-specific estimates
 of annual medical expenditures attributable to obesity. Obesity (Silver Spring). 2012
 Jan;20(1):214-20.
- Commonwealth Medicine, University of Massachusetts Medical School. MassHealth: The Basics: Facts and Trends, October 2022. Accessed September 2023. https://www.bluecrossmafoundation.org/sites/q/files/csphws2101/files/2022-10/MassHealthBasics2022_FINAL_1.pdf
- Rogers NT, Cummins S, Forde H, Jones CP, Mytton O, Rutter H, Sharp SJ, Theis D, White M, Adams J. Associations between trajectories of obesity prevalence in English primary school children and the UK soft drinks industry levy: An interrupted time series analysis of surveillance data. PLoS Med. 2023 Jan 26;20(1):e1004160. doi: 10.1371/journal. pmed.1004160. PMID: 36701272; PMCID: PMC9879401.
- Allcott H, Lockwood BB, Taubinsky D. Should We Tax Sugar-Sweetened Beverages? An Overview of Theory and Evidence. Journal of Economic Perspectives. 2019; 33 no 3: 202-27.
- Krieger J, Bleich SN, Scarmo S, Ng SW. Sugar-Sweetened Beverage Reduction Policies: Progress and Promise. Annu Rev Public Health. 2021 Apr 1;42:439-461.
- Friedman R. Public Opinion Data, 2013: New Haven, CT: Yale Rudd Center for Food Policy & Obesity.
- Federal Trade Commission. A review of food marketing to children and adolescents: follow-up report. Washington, DC Dec 2012.
- Lawman HG, Bleich SN, Yan J, LeVasseur MT, Mitra N, Roberto CA. Unemployment claims in Philadelphia one year after implementation of the sweetened beverage tax. PLoS ONE. 2019;14(3):e0213218. doi:10.1371/journal.pone.0213218
- Marinello S, Leider J, Powell LM. Employment impacts of the San Francisco sugarsweetened beverage tax 2 years after implementation. PLoS One. 2021 Jun 2;16(6):e0252094. doi: 10.1371/journal.pone.0252094

- Frieden TR, Mostashari F, Kerker BD, Miller N, Hajat A, Frankel M. Adult tobacco use levels
 after intensive tobacco control measures: New York City, 2002-2003. American Journal of
 Public Health. 2005;95(6):1016-1023.
- Colchero MA, Salgado JC, Unar-Munguia M, Molina M, Ng SW, Rivera-Dommarco JA. Changes in Prices After an Excise Tax to Sweetened Sugar Beverages Was Implemented in Mexico: Evidence from Urban Areas. PLoS One. 2015;10(12):11.
- Cawley J, Frisvold D, Hill A, Jones D. The Impact of the Philadelphia Beverage Tax on Prices and Product Availability. NBER Working Paper. 2018(24990).
 Bleich SN, Lawman HG, LeVasseur MT, et al. The association of a sweetened beverage
- Bleich SN, Lawman HG, LeVasseur MT, et al. The association of a sweetened beverage tax with changes in beverage prices and purchases at independent stores. Health Aff (Millwood). 2020;39(7):130-1139.
- Marinello S, Pipito AA, Leider J, Pugach O, Powell LM. The impact of the Oakland sugar-sweetened beverage tax on bottled soda and fountain drink prices in fast-food restaurants. Prev Med Rep. 2020;17:101034.
- Ng S, Silver L, Ryan-Ibarra S, et al. Berkeley Evaluation of Soda Tax (BEST) Study Preliminary Findings. Presentation at the annual meeting of the American Public Health Association. Paper presented at: Presentation at the annual meeting of the American Public Health Association; November, 2015; Chicago, IL.
- Powell L, Isgor z, Rimkus L, Chaloupka F. Sugar-sweetened beverage prices: Estimates from a national sample of food outlets. Chicago, IL: Bridging the Gap Program, Health Policy Center, Institute for Health Research and Policy, University of Illinois at Chicago, 2014.
- Falbe J, Thompson HR, Becker CM, Rojas N, McCulloch CE, Madsen KA. Impact of the Berkeley Excise Tax on Sugar-Sweetened Beverage Consumption. American Journal of Public Health. 2016;106(10):1865-1871.
- Cawley J, Frisvold D, Hill A, Jones D. The impact of the Philadelphia beverage tax on purchases and consumption by adults and children. Journal of health economics. 2019;67:102225.
- Lee MM, Falbe J, Schillinger D, Basu S, McCulloch CE, Madsen KA. Sugar-Sweetened Beverage Consumption 3 Years After the Berkeley, California, Sugar-Sweetened Beverage Tax. Am J Public Health. 2019;109(4):637-639.
- Zhong Y, Auchincloss AH, Lee BK, Kanter GP. The Short-Term Impacts of the Philadelphia Beverage Tax on Beverage Consumption. American journal of preventive medicine. 2018;5(1):26-36.
- Silver LD, Ng SW, Ryan-Ibarra S, et al. Changes in prices, sales, consumer spending, and beverage consumption one year after a tax on sugar-sweetened beverages in Berkeley, California, US: A before-and-after study. PLoS Med. 2017;14(4):e1002283
- California, US: A before-and-after study. PLoS Med. 2017;14(4):e1002283
 42. Lee MM, Barrett JL, Kenney EL, et al. A Sugar-Sweetened Beverage Excise Tax in California: Projected Benefits for Population Obesity and Health Equity. Am J Prev Med. 2024 Jan;66(1):94-103. doi: 10.1016/j.amepre.2023.08.004.
- Katan MB, Ruyter JC de, Kuijper LDJ, Chow CC, Hall KD, Olthof MR. Impact of Masked Replacement of Sugar-Sweetened with Sugar-Free Beverages on Body Weight Increases with Initial BMI: Secondary Analysis of Data from an 18 Month Double-Blind Trial in Children. PLoS ONE. 2016;11(7):e0159771. doi:10.1371/journal.pone.0159771
- Pan A, Malik VS, Hao T, Willett WC, Mozaffarian D, Hu FB. Changes in water and beverage intake and long-term weight changes: results from three prospective cohort studies. Int J Obes (Lond). 2013;37(10):1378-1385. doi:10.1038/ijo.2012.225
- Stern D, Middaugh N, Rice MS, et al. Changes in Sugar-Sweetened Soda Consumption, Weight, and Waist Circumference: 2-Year Cohort of Mexican Women. Am J Public Health. 2017;107(11):1801-1808. doi:10.2105/AJPH.2017.304008
 Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in Diet and Lifestyle
- Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in Diet and Lifestyle and Long-Term Weight Gain in Women and Men. New England Journal of Medicine. 201:364(25):2392-2404.
- Palmer JR, Boggs DA, Krishnan S, Hu FB, Singer M, Rosenberg L. Sugar-Sweetened Beverages and Incidence of Type 2 Diabetes Mellitus in African American Women. Archives of Internal Medicine. 2008;168(14):1487-1492.
- Schulze MB, Manson JE, Ludwig DS, et al. Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. JAMA-J Am Med Assoc.
- 2004;292(8):927-934.
 de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A trial of sugar-free or sugar-sweetened beverages and body weight in children. New England Journal of Medicine. 2012;367(15):1397-1406.
- Imamura F, O'Connor L, Ye Z, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. Br J Sports Med. 2016;50(8):496-U484.
- 51. Centers for Disease Control and Prevention. Diabetes Atlas. Accessed 10/21/23 at: https://gis.cdc.gov/grasp/diabetes/DiabetesAtlas.html#
 52. Massachusetts Executive Office of Health and Human Services. 101 CMR 314.00: Dental
- Massachusetts Executive Office of Health and Human Services. IOI CMR 314.00: Denta Services. https://www.mass.gov/doc/101-cmr-314-rates-for-dental-services/download. Published in 2018. Accessed November 16, 2020.
- 53. Ward Z, et al. NHANES III Dental Examination: An Incisive Report. unpublished report;
- Centers for Medicare and Medicaid Services. 2018 Annual EPSDT Participation Report: Form CMS-416. https://www.medicaid.gov/medicaid/benefits/early-and-periodic-screening-diagnostic-and-treatment/index.html
- Massachusetts Department of Public Health. 2018 Massachusetts Behavioral Risk Factor Surveillance System.
- Ward ZJ, Long MW, Resch SC, Giles CM, Cradock AL, Gortmaker SL. Simulation of Growth Trajectories of Childhood Obesity into Adulthood. N Eng J Med. 2017;377(22):2145-2153.
 Ward ZJ, Long MW, Resch SC, et al. Redrawing the US Obesity Landscape: Bias-Corrected
- Ward ZJ, Long MW, Resch SC, et al. Redrawing the US Obesity Landscape: Bias-Correcte Estimates of State-Specific Adult Obesity Prevalence. PLoS One. 2016;11(3):13.
 Long MW, Ward Z, Resch SC, et al. State-level estimates of childhood obesity prevalence
- Long MW, Ward Z, Resch SC, et al. State-level estimates of childhood obesity prevalence in the United States corrected for report bias. International Journal of Obesity. 2016;40(10):1523-1528.
- 59. Ward ZJ, Barrett JL, Cradock AL, Dupuis R, Lee MM, Long MW, Musicus AA, Gortmaker SL. Childhood Obesity Intervention Cost-Effectiveness Study (CHOICES) Microsimulation Model Technical Documentation: Details on Model Parameters (CHOICES v4.6.1). CHOICES Project Team at the Harvard T.H. Chan School of Public Health, Boston, MA; March 2023. Available at: https://choicesproject.org/methods/choices-model-technical-documentation

For further information, contact choicesproject@hsph.harvard.edu and visit www.choicesproject.org

APPENDIX

This appendix includes results of a \$0.02/ounce state excise tax based on the volume of sugary drinks (i.e., a flat tax).

Table A1. Results of a \$0.02/ounce state excise tax based on the volume of sugary drinks in Massachusetts, 2023-2032

Outcome	Number of people affected by the tax	95% UI
First Year Population Reach*	6.57 million	(6.55 million; 6.60 million)
10-Year Population Reach*	7.18 million	(7.14 million; 7.23 million)
Outcome	Impact of the tax on sugary drink consumption and spending	95% UI
Decrease in 12-oz Servings of Sugary Drinks per Person in the First Year*	125	(76; 230)
Decrease in Spending on Sugary Drinks in the First Year per Person Consuming Sugary Drinks*	^{\$} 82	(-\$0.04; \$255) 96% likelihood of decrease in spending
Decrease in Spending on Sugary Drinks in the First Year per Household*	\$207	(-\$0.11; \$643) 96% likelihood of decrease in spending
Decrease in Spending on Sugary Drinks in the First Year Overall in Massachusetts*	\$302 million	(-\$156,000; \$939 million) 96% likelihood of decrease in spending
Outcome	Impact of the tax on obesity and related health outcomes	95% UI
Quality Adjusted Life Years (QALYs) Gained Over 10 Years	34,300	(16,800; 73,200)
Years of Life Gained Over 10 Years	7,380	(3,020; 17,000)
Deaths Prevented Over 10 Years*	2,120	(877; 4,820)
Years with Obesity Prevented Over 10 Years	557,000	(276,000; 1.15 million)
Cases of Obesity Prevented in 2032*	72,700	(36,500; 150,000)
Cases of Childhood Obesity Prevented in 2032*	9,930	(4,090; 21,600)
Cases of Obesity Prevented Per 100,000 in 2032*	1,140	(569; 2,360)
Outcome	Impact of the tax on diabetes	
Reduction in Diabetes Incidence*	11%	
Cases of Diabetes Prevented*	2,750	

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow.

*Not discounted

Table A1. (continued) Results of a \$0.02/ounce state excise tax based on the volume of sugary drinks in Massachusetts, 2023-2032

Outcome	Impact of the tax on tooth decay	95% UI
Reduction in Decayed, Missing, or Filled Teeth among Medicaid Participants	140,000	(33,300; 141,000)
Dental Decay Treatment Cost Savings over 10 years (Medicaid)*	\$10.8 million	(\$49.4 million; \$22.1 milion)
Reduction in Decayed, Missing, or Filled Teeth among all Massachusetts residents	\$893,000	(390,000; 1.86 million)
Dental Decay Treatment Cost Savings over 10 years (Societal)*	\$68.6 million	(\$31.3 million; \$142 million)
Outcome	Costs	95% UI
Annual Implementation Cost	\$380,000	(\$328,000; \$431,000)
Annual Implementation Cost per Person	\$0.06	(\$0.05; \$0.07)
Total Intervention Implementation Cost Over 10 Years	\$3.80 million	(\$3.28 million; \$4.31 million)
Health Care Costs Saved Over 10 Years	\$1.09 billion	(\$470 million; \$2.47 billion)
Net Costs Difference Over 10 Years	-\$1.09 billion	(-\$2.46 billion; -\$466 million)
Health Care Costs Saved per \$1 Invested Over 10 Years	\$286.84	(\$123.74; \$656.77)
Outcome	Medicaid spending	95% UI
Reduction in Annual Adult Obesity-Related Medicaid Expenditures (Paid by State)*	\$3.94 million	(\$1.87 million; \$8.54 million)
Reduction in Total Annual Adult Obesity- Related Medicaid Expenditures (Paid by State & Federal)*	\$17.9 million	(\$8.52 million; \$38.8 million)
Outcome	Cost-effectiveness metrics	
Cost per Year with Obesity Prevented Over 10 Years	Cost-saving**	
Cost per Quality Adjusted Life Year (QALY) Gained Over 10 Years	Cost-saving**	
Cost per Year of Life Gained Over 10 Years	Cost-saving**	
Cost per Death Averted Over 10 Years	Cost-saving**	

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received

in the future, given that they are worth more today than they would be tomorrow. *Not discounted.
**There is a 100% likelihood that these metrics are cost-saving.

Table A2. Results of a \$0.02/ounce state excise tax based on the volume of sugary drinks in Massachusetts, 2023-2032, by race and ethnicity

	Impact of the tax on behavior and health, by race and ethnicity			
Outcome	Black or African American, not Hispanic/Latinx	Hispanic/Latinx	White , not Hispanic/Latinx	All Other Races, not Hispanic/ Latinx†
	Mean	Mean	Mean	Mean
	95% UI	95% UI	95% UI	95% UI
Decrease in 12-oz Servings of Sugary Drinks per Person in the First Year*	140 (86; 250)	133 (81; 234)	124 (75; 229)	112 (72; 211)
Quality Adjusted Life Years	2,410	4,480	24,800	2,570
(QALYs) Gained Over 10 Years	(1,190; 4,960)	(2,280; 8,830)	(12,000; 54,200)	(1,280; 5,410)
Years of Life Gained Over 10 Years	601 (190; 1,390)	592 (180; 1,390)	5,740 (2,380; 13,200)	448 (120; 1,040)
Years with Obesity Prevented	41,000	90,600	384,000	42,100
Over 10 Years	(20,700; 84,400)	(46,500; 177,000)	(188,000; 824,000)	(21,700; 82,000)
Cases of Obesity Prevented in 2032*	5,350 (2,680; 11,100)	12,000 (6,070; 23,300)	49,700 (24,500; 106,000)	5,680 (2,910; 11,100)
Cases of Childhood Obesity	1,080	3,250	4,670	925
Prevented in 2032*	(385; 2,450)	(1,270; 7,180)	(1,930; 10,200)	(332; 2,130)
Cases of Obesity Prevented Per 100,000 in 2032*	1,250 (630; 2,550)	1,360 (687; 2,620)	1,120 (551; 2,390)	909 (465; 1,780)
Comparison of Cases of Obesity	1.12	1.22	1.00 (Reference)	0.82
Prevented per 100,000 in 2032 vs. White*‡	(0.91; 1.47) 82% likelihood of greater impact	(0.95; 1.71) 94% likelihood of greater impact	N/A	(0.66; 1.05) 95% likelihood of lesser impact
	1.09	1.20	0.98	0.80
Comparison of Cases of Obesity Prevented per 100,000 in 2032	(0.93; 1.32)	(1.01; 1.50)	(0.89; 1.04)	(0.69; 0.94)
vs. Average*‡	85% likelihood of greater impact	97% likelihood of greater impact	69% likelihood of lesser impact	>99% likelihood of lesser impact

The 95% uncertainty interval (95% UI) can be thought of as a likely range. It is estimated by running the model 1,000 times, taking into account uncertainty from data sources and

population projections, and calculating a central range in which 95 percent of these model results fell.

Costs and health outcomes are discounted at 3% per year, unless otherwise noted. Discounting estimates the present value of costs and health outcomes that are spent or received in the future, given that they are worth more today than they would be tomorrow. *Not discounted.

[†] All Other Races includes people of Asian. Native Hawaiian and Pacific Islander, American Indian and Alaska Native, and two or more races,

[‡] The ratio of the projected reduction in obesity prevalence in a particular group compared to the projected reduction in the reference group. A value greater than 1 indicates that the group is projected to see a greater reduction in obesity levels compared to the reference group.